



**International Journal of Biology, Pharmacy  
and Allied Sciences (IJBPAS)**

*'A Bridge Between Laboratory and Reader'*

[www.ijbpas.com](http://www.ijbpas.com)

---

---

**STUDY OF THE DISTRIBUTION OF THE MOST COMMON GENETIC DISEASES IN  
THE LAST FIVE YEARS BY USING GIS (CASE STUDY: KURDISTAN PROVINCE)**

**DR. ASSADOLAH FATHOLAH POOR<sup>1</sup>, GOLALEH KARBASSI<sup>2</sup>, SHLER  
KATORANI<sup>3</sup>, KAJEJEH KARBASI<sup>4</sup>**

1-Assistant Professor of Pediatrics Department, Faculty of Medicine, Kurdistan University of  
Medical Sciences, Sanandaj, Iran, [Gelare\\_633@yahoo.com](mailto:Gelare_633@yahoo.com)

2-Master of Remote Sensing and GIS, Center for Community Services gis Expert , Jihad  
University of Kurdistan, Kurdistan, Iran, [sh.katorani@yahoo.com](mailto:sh.katorani@yahoo.com)

3-Master of pediatric Nursing, Genetics expert Kurdistan welfare, Iran (Corresponding author)

4- Bachelor in Natural Sciences, Iran

**ABSTRACT**

Identification of the accumulation of certain diseases in the province and the country will considerably be helpful for the human resources management, allocation of budget, and health services in centers of accumulation of the diseases. Considering that so far no project has been conducted on the issue of spatial analysis of genetic disease in the Kurdistan province, preparing the maps of prevalence of genetic diseases, determining the geographical direction of the spread of these disease in the province of Kurdistan and specifying the areas susceptible to disease and centers of accumulation of genetic disorders in Kurdistan province will significantly be helpful in identifying the way that geographic areas are influential in the creation of these diseases. According to the graphs and tables, it was found that the rate of the prevalence of phenylketonuria, in Kurdistan Province, is higher in males than in females and the highest number of disease were seen in children aged between 1 and 5 years old and the rate of prevalence of this disease in Saqqez is higher than other cities of Kurdistan province. Also, in Kurdistan Province, the rate of prevalence of Thalassemia is higher among males than females and the greatest numbers of patients were aged between 21 to 25 years old and according to

---

statistics the rate of prevalence of this disease in Sanandaj, is higher than other cities of Kurdistan province. Also the rate of prevalence of hemophilia, in Kurdistan, is higher in males than the females and most of the patients were aged between 11 to 30 years old, and figures show that the highest rate of prevalence of the disease was seen in Sanandaj. The prevalence rate of physical motor impairment with genetic causes, in Kurdistan, is more in males than females and it has been reported that the statistics of the prevalence of this disease in Bijar, is higher than the other cities of Kurdistan. Also the prevalence of blindness and mental disabilities with genetic causes, in Kurdistan, is more in males than females and the rate of the prevalence of this disease in Sanandaj, is higher than the other cities of Kurdistan and the prevalence of hearing loss and deafness with genetic causes, in Kurdistan, is more in males than females and the rate of the prevalence of this disease in Kamyaran, is higher than the other cities of Kurdistan

**Keywords: GIS, genetic disease, Geographical distribution**

## **1- INTRODUCTION**

Geographic information system has an efficient and significant role in monitoring and evaluation of diseases and in the field of hygiene & health. By using this knowledge, the areas of accumulation of the diseases can be placed at the disposal of the authorities of healthcare centers in the form of graphs, maps and tables. Spatial analysis of various diseases, with the use of geographic information system, indicates the distribution pattern of diseases in different regions of the earth that this issue leads to the examination of the causes of its distribution in order to find that whether the causes of the distribution are environmental or weather and climatic factors or that other problems have caused the distribution of the disease with their specific pattern. If the disease is

concentrated in certain parts of the study area, then the causes of this matter should be searched in these regions and the type of food, water and weather and also eating habits of these regions and every factor that exists in that region and can be helpful in the creation and spread of the disease should be investigated. One of the most important diseases that affect human is genetic diseases that in terms of treatment and recovery it imposes the greatest costs on the patient and after spending high costs, the chance of recovery of the patient is very low. Many of genetic diseases are incurable and high costs will be imposed at the stages of preventing the spread and the development of disease in the patient prone to these disease. In intra-familial marriage, the possibility of incidence

of the disease is much higher, for this reason the genetic counseling and genetic testing for avoiding the birth of children with genetic diseases are among the most important ways to prevent individuals from being affected by these diseases. Identification of the accumulation of certain diseases in the province and the country will considerably be helpful for the human resources management, allocation of budget, and health services in centers of accumulation of the diseases. Considering that so far no project has been conducted on the issue of spatial analysis of genetic disease in the Kurdistan province, preparing the maps of prevalence of genetic diseases, determining the geographical direction of the spread of these disease in the province of Kurdistan and specifying the areas susceptible to disease and centers of accumulation of genetic disorders in Kurdistan province and also determination of the coefficient of influence of climatic and environmental parameters in the development of genetic diseases with respect to the geographical, climatic and situational parameters of the study area will significantly be helpful in identifying the way that geographic areas are influential in the creation of these diseases. Studying the spatial factors that are involved in the development of the disease seems to be

necessary. In the present study, it has been tried to investigate the distribution pattern of genetic disease in Kurdistan province, and to model the causes of this distribution and effective parameters.


## **2- MATERIALS AND METHODS**

### **2-1 The study area**

Kurdistan province with an area of approximately 28,203 square kilometers is one of the western provinces of Iran that has been placed in neighborhood of the provinces of West Azerbaijan, Zanjan, Hamedan and Kermanshah and it is on the border with Iraq and there are more than 200 kilometers of common border with Iraq. The geographical coordinates of the Kurdistan Province are at 34 degrees 44 minutes up to 36 degrees 30 minutes north latitude and 45 degrees 31 minute to 48 degrees and 16 minutes eastern longitude. Sanandaj is the capital of Kurdistan Province and is located at 1373 meters above the sea level. Other cities of the province are: Saqqez, Marivan, Qurveh, Baneh, Bijar, Dehgolan, Divandarreh, Kamyaran and Sarvabad. This province is located on the scattered slopes and plains of central Zagros Mountain and it is bound by Iraq on the west, the province of West Azerbaijan and Zanjan to its north, Zanjan and Hamedan to the east and Kermanshah to the south. Kurdistan Province is entirely a

mountainous region and roughness of the province consists of two western and eastern sections. Wide range of Sanandaj, Marivan and its surrounding lands up to the south of the Kurdistan constitute the western roughness. East of the Sanandaj is covered by the eastern roughness. This area is

surrounded with the mountainous fence that is composed of metamorphic and sedimentary rocks. The highest mountains of this region, Shah Neshin, are located in the north of Bijar, Sheida at the center and Panjeh Ali between Qurveh and Saqqez (Figure 1).



**Figure 1- Position of Kurdistan province in Iran**

According to the latest political and administrative division of 2006, Kurdistan Province has 9 cities, 26 counties, 84 districts and 1864 villages that 1765 villages are inhabited and 99 villages are haunted. Cities of Kurdistan province are: Saqqez, Qurveh, Bijar, Sanandaj, Marivan, Baneh, Divandarreh, Kamyaran, Sarvabad that Sanandaj is the capital of Kurdistan province.

### 3- Examined variables

Data of the disease include ID card specifications and medical records of the patients in Kurdistan Province, from 2009 until the end of 2014 and they have been prepared and gathered from the Welfare Organization and health care center of University of Medical Sciences of Kurdistan province. This information includes personal information such as age, gender and address of patients and specifications related to

medical diagnosis and the type of disease. All of these specifications were collected for the diseases of hemophilia, phenylketonuria, thalassemia, physical motor disability, mental disability, blindness and deafness with the genetic causes.

#### 4- CONCEPTUAL MODEL OF THE RESEARCH

In the first stage, the required spatial data were collected (such as maps of cities and villages, political and administrative boundaries of provinces, cities and counties, etc.) The spatial data were prepared and collected from map preparing agencies such as the National Cartographic Center and Cadaster of the country and other organizations such as the General Governor Office, Housing and Urban Development, Housing Foundation and ....

In the second stage, the demographic data and maps of the province, cities, districts and

the province's political divisions were prepared.

In the third stage, the data of the disease were collected that this task was carried out by referring to welfare centers or health care centers of University of Medical Sciences.

In the fourth stage, the relevant software for the data analysis were determined that included ARC GIS and EDRI SI software and methods of spatial analysis and modeling techniques were identified.

In the fifth stage, spatial and statistical analyses were applied on disease and the maps of prevalence and spread of disease were prepared and the type of distribution and the direction of the spread of disease in the province were specified. In the last stage the results were displayed in the form of maps, tables and graphs.




Figure 2- Conceptual model of the research


In order to prepare a map of the geographical distribution of disease in Kurdistan province, initially, the spatial maps of the province including maps of the province, city, district and village were placed in a GIS environment and then a data layer was prepared in GIS environment with the geometric point in order to enter the data about the patients. According to the data collected from patients in Kurdistan province in the statistical years of 2009 until the end of 2014 and with regard to the exact address of patients, for every patient a point was considered in the position of the patient's address and consequently the stages of preparation of the map was completed and the output was a map which included borders





of the province, borders of cities and areas of disease. All the spatial - statistical analysis will be carried out on the basis of this basic map.

In the following part of the research, the map of the geographical distribution of the disease is drawn for the most common genetic disease in the province in the statistical years 2009 to 2014 and according to the number of patients who have referred to genetic counseling centers in Welfare Organization of the province in the border of cities of Kurdistan province.

**5. The spatial analysis of disease**

In order to determine the direction of spread of disease, a standard deviation ellipse and mean center is drawn.








Figure 3- Geographical distribution and standard deviation ellipse of diseases in Kurdistan province

## 6- FINDINGS

### 6-1 The situation of phenylketonuria (PKU) in Kurdistan Province

Considering the information obtained from the Welfare Organization and health care center of University of Medical Sciences of Kurdistan province, the number of patients with this disease, according to their gender, age and city of residence is as follows.


### 6-2 The situation of thalassemia in Kurdistan Province

Considering the information obtained from the Welfare Organization and health care center of University of Medical Sciences of

Kurdistan province, the number of patients with this disease, according to their gender, age and city of residence is as follows.

### 6-3 The situation of hemophilia in Kurdistan Province

Considering the information obtained from the Welfare Organization and health care center of University of Medical Sciences of Kurdistan province, the number of patients with this disease, according to their gender, age and city of residence is as follows.




Gender	Frequency
Male	34
Female	29
Total	63


Table and Figure 1- Data of Phenylketonuria disease according to the gender of patients in Kurdistan Province (2014)

Table 2- Data of Phenylketonuria disease according to the age of patients in Kurdistan Province (2014)

Age (Year)	1 to 5	6 to 10	11 to 15	16 to 20	21 to 25	26 to 30	31 and more
Frequency	31	9	9	2	4	3	5



Graph 3-Data of Phenylketonuria disease according to the age of patients in Kurdistan Province (2014)



Graph 4- Distribution of Phenylketonuria in the cities of Kurdistan Province (2014)






Table and Figure 5- Data of thalassemia according to the gender of patients in Kurdistan Province (2014)

Table 6-Data of thalassemia disease according to the age of patients in Kurdistan Province (2014)

Age (Year)	1 to 5	6 to 10	11 to 15	16 to 20	21 to 25	26 to 30	31 and more
Frequency	6	15	32	29	23	21	4



Graph 7-Data of thalassemia disease according to the age of patients in Kurdistan Province (2014)



Graph 8- Distribution of thalassemia in the cities of Kurdistan Province (2014)






Table and Figure 9- Data of hemophilia according to the gender of patients in Kurdistan Province (2014)

Table 10- Data of hemophilia disease according to the age of patients in Kurdistan Province (2014)

Age (Year)	6 month to 10	11 to 30	31 to 50
Frequency	15	75	32



Graph 11- Data of hemophilia disease according to the age of patients in Kurdistan Province (2014)



Graph 12- Distribution of hemophilia in the cities of Kurdistan Province (2014)

**6-4 The situation of mental disabilities in Kurdistan Province**

Considering the information obtained from the Welfare Organization and health care center of University of Medical Sciences of Kurdistan province, the number of patients with this disease, according to their gender, age and city of residence is as follows.


**6-5 The situation of hearing loss and deafness in Kurdistan Province**

Considering the information obtained from the Welfare Organization and health care

center of University of Medical Sciences of Kurdistan province, the number of patients with this disease, according to their gender, age and city of residence is as follows.


**6-6 The situation of blindness in Kurdistan Province**

Considering the information obtained from the Welfare Organization and health care center of University of Medical Sciences of Kurdistan province, the number of patients with this disease, according to their gender, age and city of residence is as follows.




Gender	Frequency
Male	314
Female	192
Total	506

Table and Figure 13- Data of mental disabilities according to the gender of patients in Kurdistan Province (2014)



Graph 14- Distribution of mental disabilities in the cities of Kurdistan Province (2014)



Gender	Frequency
Male	298
Female	237
Total	535

Table and Figure 15- Data of hearing loss and deafness according to the gender of patients in Kurdistan Province (2014)

Graph 12- Distribution of hearing loss and deafness in the cities of Kurdistan Province (2014)






Table and Figure 17- Data of blindness according to the gender of patients in Kurdistan Province (2014)



Gender	Frequency
Male	114
Female	89
Total	203




Graph 18- Distribution of blindness in the cities of Kurdistan Province (2014)

**6-7- The situation of physical disabilities in Kurdistan Province**


Considering the information obtained from the Welfare Organization and health care

center of University of Medical Sciences of Kurdistan province, the number of patients with this disease, according to their gender, age and city of residence is as follows:



Gender	Frequency
Male	290
Female	180
Total	475

Table and Figure 19- Data of physical disabilities according to the gender of patients in Kurdistan Province (2014)



Graph 20- Distribution of physical motor disabilities in the cities of Kurdistan Province (2014)

**7- Comparison of the prevalence of the studied diseases in the province and their prevalence in the country:**

According to the graphs and tables, it was found that the rate of the prevalence of phenylketonuria, in Kurdistan Province, is higher in males than in females and the highest number of disease were seen in children aged between 1 and 5 years old and the rate of prevalence of this disease in Saqqez is higher than other cities of

Kurdistan province. Also, in Kurdistan Province, the rate of prevalence of Thalassemia is higher among males than females and the greatest numbers of patients were aged between 21 to 25 years old and according to statistics the rate of prevalence of this disease in Sanandaj, is higher than other cities of Kurdistan province. Also the rate of prevalence of hemophilia, in Kurdistan, is higher in males than the females and most of the patients were aged

between 11 to 30 years old, and figures show that the highest rate of prevalence of the disease was seen in Sanandaj. The prevalence rate of physical motor impairment with genetic causes, in Kurdistan, is more in males than females and it has been reported that the statistics of the prevalence of this disease in Bijar, is higher than the other cities of Kurdistan. Also the prevalence of blindness and mental disabilities with genetic causes, in Kurdistan, is more in males than females and the rate of the prevalence of this disease in Sanandaj, is higher than the other cities of Kurdistan and the prevalence of hearing loss and deafness with genetic causes, in Kurdistan, is more in males than females and the rate of the prevalence of this disease in Kamyaran, is higher than the other cities of Kurdistan. According to the monitoring and evaluation of some genetic diseases, including thalassemia, hemophilia and phenylketonuria (PKU), which have been conducted in the country and considering the reports obtained from the center of the health care of Medical Sciences Universities of the country and also Kurdistan province and also the studies that have been performed in the present research, it was found that the prevalence of the mentioned genetic disorders in Kurdistan province is less than other provinces of the

country and in the state divisions, in terms of the prevalence of genetic disorders, Kurdistan province is placed in the position of mild to moderate and northern provinces, in terms of the prevalence of genetic disorders, are ranked first in the country and according to the state divisions they are placed in the category of severe to moderate. However, in the case of disease of deafness, according to conducted surveys and statistics and data available in the Welfare Organization, the latest reports indicate that the country's average of deafness disease is 2.6 that this figure is 2.2 in Kurdistan province and this represents that the index of deafness of Kurdistan Province is close to the national average and hence, the prevalence of deafness in Kurdistan province is located in the middle class and this matter can be a wake-up call for the authorities of Kurdistan Province to investigate the cause of this issue. Many of the diseases impose high financial costs on patients and their families. Investigating the causes of the disease and the prevalence of a number of diseases in the province and examining the status of the disease in the country requires laboratory studies and also the identification of the level of impact of all factors related to disease that as a result of it the factor or factors of the prevalence of intended diseases will be

detected in the province or country and the necessary measures would be carried out to reduce or eliminate these factors. Identification of the accumulation of certain diseases in the province and the country will considerably be helpful for the human resources management, allocation of budget, and health services in centers of accumulation of the diseases.

## 8- REFERENCES

- [1] Aykhaee Bijan, Zandian Khodamorad (2004). Reporting the new findings, in order to revise the way of prevention of thalassemia major and sickle cell anemia in Iran. *Medical Science Journal*, 77 to 85.
- [2] Behbehani, S. Karimi Maridani, M (2010). Neural network and its application in medical engineering, *Journal of Medical Engineering*, No. 115, 41-54
- [3] Behrouzi Khazaei, Kh. Tavakkoli, T. Amiri Chaijan, R. Khosh Tagaza, M. (2010). Application of artificial neural network to predict the moisture content during the process of drying grapes, *Proceedings of the Fifth National Congress of Agricultural Engineering and Mechanization*, Ferdowsi University of Mashhad.
- [4] Pakzad, M; *Exceptional Children*, Tehran, ISBN, 1997, third edition, p. 345.
- [5] Pezeshki, Z. Zahrayi, M. Tafazzoli Shadpour, M. Omid, A. (2009). The incidence of cholera and its relationship with climatic conditions during the years 1995-2006, *Journal of Infectious and Tropical Diseases affiliated with the Association of Tropical and Infectious Diseases Specialists*, No. 45, pages 11-15.
- [6] PourFathollah Ali Akbar, Rahgozar Soheila, Maafi Alireza (2003) "The importance of an accurate measurement of hemoglobin F in the differential diagnosis of sickle cell anemia of sickle beta thalassemia, *Journal of Medical Council of Islamic Republic of Iran* 51-54.
- [7] Pirmoradi, A. Nouri fard, M. Salahi mogaddam, A. (2012). Ecological study of malaria in the city of Bandar Abbas, by using Geographic Information Systems, *Journal of Military Medicine of the Islamic Republic of Iran*, No. 1, pages 35 -44.
- [8] Hossein Zadeh, M. Servati, M. Mansouri, A. Mirbagheri, B. And khezri, C. (2009). Zoning the risk of incidence of mass movements by

- using logistic regression model (case study of the Sanandaj - Dehgolan), *Journal of Geology of Iran*, Third Year, Issue 11, pages 27-37.
- [9] khalili, A. Darvish Sefat, A. Baradaran Rad, R. Bazr Afshan, G. (2004)..A suggestion for climate classification in GIS, Case study: northwest of Iran, *Desert Journal*, No 2.
- [10] Khoshdel, A. Nouri Fard, M. Pezeshkan, R. Salahi Mogaddam, A. (2012), Mapping of contagious diseases of Iran, the *Journal of Health and Development*, No. 1.
- [11] Rahmani, A. Esmaili, GH. (2010).The Efficiency of Neural Networks, Logistic Regression & Discriminant Analysis in Defaults Prediction, *Journal of Quantitative Economics (former economic studies)*, No. 4, pages 152-172.
- [12] Zare, M. Shams Zadeh, P. Najjari, A. (2005).Application of GIS in improving decisions of health sector management, *Journal of University of Medical Sciences of Gilan*, No. 59.
- [13] Sadhi, M. Mehrabi, Y. and Khodabakhshi, A. (2011).The use of principal component analysis to increase the accuracy of prediction of the metabolic syndrome in the artificial neural network and logistic regression models, *University of Medical Sciences*, No. 4, pages 18-27.
- [14] Sayyadi, A. (2008).A preliminary familiarization with artificial neural networks, laboratory of processing vital signals and medical images, Faculty of Electrical Engineering, Sharif University.
- [15] Katorani, Sh. Almodarresi, A. Karami, G. 2010.Geographical distribution study of skin cancer cases in Kurdistan province, a Master's thesis.
- [16] Kakavand, A; 2006,Psychology and Education of Exceptional Children, University of Social Welfare and Rehabilitation Sciences, Tehran,First Edition, p. 178.
- [17] Kazemi, A; Karimlo, M; Rahgozar, M; Bakhshi, A. A; Asgari, A. (2011),Application of Bayesian method in the estimation of the parameters of the logistic regression model with random missing values of auxiliary variables, North Khorasan University of Medical

- Sciences, Special Issue - Biostatistics and Epidemiology.
- [18] Kiani, B. Tabari, M. Fallah, A. Hosseini, M. and Iran Nezhad, H. (2011).The use of three methods of: nearest neighbor,Ripley's K function and mean square to determine spatial pattern of Saxaul (Haloxylon) in Siahkooh protected area, Yazd province, Iranian Journal of Forest and Poplar Research, Issue 3, Pages 356-369.
- [19] Mesgari, M. Masoumi, Z. (2005).Using statistical and spatial analysis of the geographic information systems in the discovery of spatial concentration of diseases (case study: Discovering the spatial concentration of gastric cancer in Iran), Geomatics Conference
- [20] Mozaffari, Gh. Bakhshi Zadeh Kloucheh, F. Gheibi, M. (2010). Investigating the relationship between the vegetation cover and prevalence rate of cutaneous leishmaniasis by using Geographical Information System, Case Study: Ardekan- Yazd, Journal of Geography and Environmental Planning, issue 44, pages 167- 178.
- [21] Masoumi, Z. Mesgari, M. (2006).Fitting of a spatial- statistical model on skin cancer to support the decision making, Geomatics Conference, 2006,National Cartographic Center, Tehran, of Iran.
- [22] Masoumi, Z. Mesgari, M. Karimi, M. (2010).Simulation of spatial distribution of skin cancer by using statistical-spatial analysis, Journal of Civil Engineering and Mapping, Faculty of Engineering, No. 5, pages 753-763.
- [23] Mirzadeh, A. Hejazi Zadeh, B. Mesgar pour, B. Golezar, A. Halakooiiy Nayini, K. (2008), The process of development of high-risk centers of cutaneous leishmaniasis in the city of Kerman in 2002-2005 years andevaluation of effective environmental factors by using GIS, Iranian Journal of Epidemiology, Vol. 4, No. 3 and 4, pages 17-5.
- [24] Anselin.L,) 2003(. An Introduction to Point Pattern Analysis using CrimeStat, Spatial Analysis Laboratory, Department of Agricultural and Consumer Economics, University of Illinois, Urbana-Champaign.

- [25] Antshel KM. (2010). ADHD, learning, and academic performance in phenylketonuria. *Mol Genet Metab.* 99 Suppl 1:S52-8.
- [26] Berntorp E. 2009 Joint outcomes in patients with haemophilia: the importance of adherence to preventive regimens. *Haemophilia*; 15(6): 1219- 27.
- [27] Brumm VL, Grant ML) 2010(. The role of intelligence in phenylketonuria: a review of research and management. *Mol Genet Metab.*;99 Suppl 1:S18-21
- [28] Cockings, S. Bhopal,c. Walker,d,D. (2004). Users' perspectives on epidemiological,GIS and point pattern approaches to analysing environment and health data, *Health & Place*, 10, PP 169–182.
- [29] Cressie N.A, 1993. *Statistics for spatial data*, Wiley series in Probability and Mathematical Statistics. P900.
- [30] Cromley, E. and McLafferty S)2002(. *GIS and public health*. New York: The Guilford Press.
- [31] Dayhoff, J. E.1990. *Neural Network Principles*, Prentice-Hall International, U.S.A.
- [32] Deane W. (2001). Use of a density equalizing map projection in analyzing childhood cancer in four California counties, *statistics in medicine*, 20,pp 1499–1513.
- [33] Demierre MF, Maguire-Eisen M, O'Donnell N, (2009). A sun protection community intervention in Quincy middle schools: Insights from the use of ultraviolet photography and its impact on sunburn. *Journal of the Dermatology Nurses' Association*,1,PP.111-18.
- [34] Dietary information for the treatment of Phenylketonuria (2008). The National Society for Phenylketonuria (United Kingdom) limited (NSPKU). Available at: Access dat: 18.
- [35] Di Nocera. F, Terenzi. M, Camilli. M,(2006). Another look at scanpath: distance to nearest neighbor as a measure of mental workload. Maastricht, the Netherlands: Shaker Publishing. pp. 1 - 9.
- [36] Getis, A. Ord, J. K. (1992). The analysis of Spatial Association by use of Distance Statistics. *Geographic Analysis*, Vol. 24, No. 3, pp. 189-206.

- [37] Gloster HM, Brodland DG. (1996). The Epidemiology of sKIN Cancer. *Dermatol Surg*; 22:PP, 217-26.
- [38] Ghiasvand NM, Aledavood A, Ghiasvand R, Seyedin Borojeny F, Aledavood AR, Seyed S, *et al.* 2009. Prevalence of classical phenylketonuria in mentally retarded individuals in iran. *J inherits metab dis.* 32 suppl., 1:S283-7.
- [39] Janzen D, Nguyen M. (2010), Beyond executive function: non-executive cognitive abilities in individuals with PKU. *Mol Genet Metab.* 99 Suppl 1:S47-51.
- [40] Kassandra,A. Matthew, K. Rebecca, B. (2009). Use of GIS to identify optimal settings for cancer prevention and control in African & American communities, *Preventive Medicine*, 49,pp54–57.
- [41] Khanna, T. (1990). *Foundation of neural networks.* Addison-Wesley Publishing Company, U.S.A.
- [42] Khawaji M, Astermark J, Von Mackensen S, Akesson K, Berntorp E. 2011 )Bone density and health-related quality of life in adult patients with severe haemophilia. *Haemophilia* 17(2): 304-11
- [43] Kulldorff, Martin, and Neville Nagarwalla. (1995). spatial disease clusters: Detection and inference. *Statistics in Medicine*, 14, PP 799-810.
- [44] Lai.P.C, Mun So.F, Wing.C.K, (2008). *Spatial epidemiological approaches in disease mapping and analysis.* Baco Raton, CRC Press.
- [45] Leshin B, White WL. (1996) malignant neoplasm's of keratinocytes. In Arndt KA, Leboi. PP 440-1378.
- [46] Levine.N, (2010). *Spatial Centographic & Nearest Neighbor Measures an Introduction to CrimeStat III,* th National Crime Mapping Conference Manchester.
- [47] Little PF. Structure and function of the human genome. *Genome Res.* 2005) Dec; 15(12):1759-66.
- [48] MESGARI, M.S. MASOOMI, Z. (2008). GIS Applications in Public Health as a Decision Making Support System and It's Limitation in Iran, *WORLD APPLIED SCIENCES JOURNAL, SUPPLE* 1,PP. 73-77
- [49] Mosavi-Jarrahia, A. Moini, M. Mohagheghi, M. Alebouyeh, M. Yazdizadeh, B. Shahabian, A.

- Nahvijo, A. Alizadeh, R. (2007). Clustering Of Childhood Cancer In The Inner City Of Tehran Metropolitan Area: A Gis-Based Analysis, International Journal Of Hygiene And Environmental Health, 210, PP113-119.
- [50] Ord, J. K., and Getis, A. (1995). Local Spatial Autocorrelation Statistics, Distributional Issues and an Application. Geographic Analysis, Vol. 27, No. 4, pp. 286-306.
- [51] Robinson, G. M. (1998). Methods and techniques in human geography. Chichester: John Wiley & Sons.
- [52] Ross C, Goldenberg NA, Hund D, Manco-Johnson MJ.( 2009) Athletic participation in severe hemophilia: bleeding and joint outcomes in children on prophylaxis. Pediatrics; 124(5): 1267-72
- [53] Tuncbilek E, Ozguc M.( 2007 )Application of medical genetics in Turkey. Turk J Pediatr. 49(4): 353-9.
- [54] Williams RA, Mamotte CD, Burnett JR. Phenylketonuria (2008).An inborn error of phenylalanine metabolism. Clin Biochem Rev. 29(1):31-41.
- [55] Yesilipek M. (2007)) Stem cell transplantation in hemoglobinopathies. hemoglobin; 31: 251-6.