

**EFFECT OF MANGANESE IN RELATION TO ANTIOXIDATIVE METABOLISM IN
GERMINATING MUNG BEANS**

***ANURADHA GOYAL¹ AND SUMIT CHHIBBER²**

¹Department of Biotechnology, S.D. College, Ambala Cantt

²Department of Botany, S.D. College, Ambala Cantt

***Corresponding Authors: E Mail: mittalanuradha7@gmail.com**

ABSTRACT

During past few decades, heavy metals, constitute important groups of environmental pollutants, have received increasing attention. Heavy metals are non-degradable which enter the food chain and subsequently to human and animal health. In humans, it can cause neurodegenerative diseases and other symptoms like, increased allergic reactions, high blood pressure, depression, irritability, poor concentration, sleep disabilities etc., some of the many conditions resulting from exposure to toxins. In plants, metal toxicity is manifested through an array of physiological and metabolic alterations including altered activities of key enzymes, synthesis of metal detoxifying compounds and due to induction of oxidative stress. Most of metals when accumulate in plant tissues cause increased generation of reactive oxygen species. In organisms, antioxidants help to deal with oxidative stress caused by free radical damage; as the main characteristic feature of an antioxidant is its ability to trap free radicals. The present work was done to investigate the activity of antioxidative enzymes in germinating mung beans under varying concentration of manganese. The germination of seeds depends on the concentration of manganese metal, under low concentrations; a considerable amount of growth is shown as it is required by the plant in trace amount. But as the concentration increases the growth of plant is inhibited due to free radical produced during stress condition of manganese. Trace elements are necessary for the normal metabolic function of the plants, but at higher concentrations, these metals are toxic and may severely interfere with physiological and biochemical functions.

Keywords: Antioxidative enzyme, metal stress, mung beans, germination

INTRODUCTION

In recent years heavy metal pollution has become one of the serious environmental problems worldwide. Unlike organic pollutants that can be easily degraded to harmless small molecules, toxic elements, such as lead, mercury, cadmium, copper and zinc, are immutable by biochemical reactions [1] hence, it is difficult to remediate these metals from the soil and water. The hot spots of soil contamination are located in the regions of large industrial activities, where surrounding agricultural lands are affected by the deposition of heavy metals and also agricultural practice, e.g., application of sewage sludges, phosphate fertilizers, liming, irrigation water and pesticides has lead to increased heavy metal concentration in soils [2]. The toxicity of metals can affect functioning of all type of living organism from microorganism to higher organisms.

In humans, it can cause neurodegenerative diseases and other symptoms like, increased allergic reactions, high blood pressure, depression, mood swings, irritability, poor concentration, aggressive behavior; sleep disabilities, fatigue, speech disorders, high blood pressure, neuropathy, autoimmune diseases, and chronic fatigue are just some of the many conditions resulting from exposure to toxins.

Very harmful thing that is produced by metal toxicity is production of free radical. When cells in the body encounter a free radical, the reactive radical may cause destruction in the cell. Oxygen free radicals induce damage due to peroxidation to biomembranes and also to DNA which lead to tissue damage, thus cause occurrence of number diseases. About 5% or more of the inhaled oxygen is converted to reactive oxygen species (ROS) such as O_2^- , H_2O_2 and OH^- by univalent reduction of O_2 . With the presence of metal ion pollution the synergistic effect of metals and ROS is dangerous to body.

Antioxidant is any substance that, when present at low concentrations, significantly delays or prevents oxidations of cell contents like proteins, lipids, carbohydrates and DNA. Antioxidants neutralize the effect of free radicals through different ways and may prevent the body from various diseases [3]. Anti oxidants can act by scavenging reactive oxygen species like Superoxide Dismutase (SOD) removes O_2^- , by inhibiting their formation i.e. by blocking activation of phagocytes, by binding transition metals ions and preventing formation of OH^- or decomposition of lipid hydroperoxides, by repairing damage. Some of the antioxidant enzymes that are found to

provide protection against the ROS are superoxide dismutase, catalase, peroxidase, and ascorbate peroxidase [4].

Mung bean is a very popular pulse crop in Asian countries, due to which it was selected for the present study. The current research was an attempt to understand the effect of Mn on growth and on antioxidant enzymes of mung bean.

MATERIALS AND METHODS

Plant material and seed collection

A cultivar of Mung bean (*Vigna radiata* L.) was used as the source material in the present investigation. The seeds of the cultivar were obtained from Ambala, Haryana (India).

Germination of seeds

Mung bean seeds were taken and washed with distilled water. After that the seeds were surface sterilized with 0.1% HgCl₂ for 5 min and rinsed with sterile distilled water for 5-6 times. Surface sterilized seeds were germinated aseptically in petri plates having sterilized wet filter paper. The experiments were performed in duplicate.

Extraction of enzymes

Enzymes were extracted by homogenizing germinated seeds (1 g) in 10 ml of the extraction buffer (50mM potassium phosphate buffer, pH 7.0) using a chilled mortar and pestle. The resulting homogenate was centrifuged at 10,000 x g for 15 min at

4°C and the supernatant was used for the determination of activities.

Estimation of catalase (CAT) activity

Catalase (EC 1.11.1.6) activity was determined [5]. The assay mixture for determining CAT activity containing 1.4 ml potassium phosphate buffer (50mM, pH 7.0), 1.5 ml of H₂O₂ (12.5 mM) and 0.1 ml of enzyme extract was used to give a final reaction volume of 3 ml. The reaction was started by the addition of H₂O₂. A blank was run without addition of the enzyme extract. The decrease in H₂O₂ was followed by recording the decrease in absorbance at 240 nm for 3 min at 30s intervals in a double beam UV-VIS spectrophotometer. One unit (U) of CAT activity was defined as the amount of enzyme catalyzing the decomposition of 1 μmol H₂O₂ per min at 240 nm.

Estimation of ascorbate peroxidase (APX) activity

Ascorbate peroxidase (EC 1.11.1.11) activity was determined [6]. The 3.0 ml of assay mixture containing 1.0 ml potassium phosphate buffer (50 mM, pH 7.0), 1.0 ml H₂O₂ (39 mM), 0.8 ml ascorbic acid (0.5 mM) and 0.2 ml of the enzyme extract. The reaction was started by the addition of H₂O₂. A blank was run without addition of the enzyme extract. The activity of APX was

measured by monitoring the rate of ascorbate oxidation at 290 nm for 3 min at 30s intervals in a double beam UV-VIS spectrophotometer. One unit of APX activity was defined as the amount required to decompose 1 μ mol ascorbic acid oxidized min⁻¹ calculated from the extinction coefficient of 2.6 mM⁻¹ cm⁻¹.

Estimation of peroxidase (POX) activity

Peroxidase (EC 1.11.1.7) activity was assayed [7]. The assay mixture comprised o-dianisidine (2.4 μ mol), H₂O₂ (20 μ mol), crude extract (0.05-0.5 mg protein) and 0.05 M citrate buffer (pH 4.8) to make the final volume of reaction mixture 3 ml and omission of H₂O₂ from the incubation mixture served as a blank. The enzyme activity was measured by following the absorbance at 430 nm at intervals of 15 sec. One unit of enzyme activity represents a change of 1 unit/min of absorbance at 430 nm.

Estimation of H₂O₂

The amount of hydrogen peroxide was estimated [8]. Supernatant was diluted to 2 ml with 10 mM potassium phosphate buffer (pH.7.0). 2ml 5% potassium dichromate and glacial acetic acid (1:3 v/v) was added to the reaction mixture. The absorbance was read at 570 nm against the reagent blank without sample extract .The quantity of H₂O₂ was

determined from the standard curve prepared by taking different concentrations of H₂O₂ ranging from 20 to 100 moles.

Estimation of malondialdehyde (MDA)

Malondialdehyde (MDA) was estimated [9].To 1 ml of supernatant was added an equal volume of MDA reagent (20% TCA in 5% thiobarbituric acid) and kept in a water bath at 95°C for 40 min and immediately chilled on ice for 15 min. The mixture was centrifuged at 10,000 x g for 30 min and the absorbance of the supernatant was measured at 520 nm and 600 nm. The non-specific absorbance at 600 nm was subtracted from that of 520 nm. The content of MDA was calculated using the extinction coefficient 155 mM⁻¹ cm⁻¹.

Estimation of Superoxide Dismutase (SOD)

Superoxide Dismutase (SOD) was estimated using pyragallol (5% v/v). To 0.5 ml of 0.1 M Tris-HCl (pH 8.2), equal volume of distilled water, pyragallol and enzyme extract was added to give a final reaction volume of 2 ml. The activity of SOD was measured by monitoring the autooxidation of pyragallol at 420 nm for 3 min at 30s intervals in a double beam UV-VIS spectrophotometer. One unit of enzyme activity represents a change of 1 unit/min of absorbance at 420 nm.

Estimation of soluble proteins

Soluble proteins were estimated [10]. The amount of soluble proteins was calculated in mg g⁻¹ FW with the help of standard plot of BSA (0-150 µg).

Statistical Analysis

All the experiments were performed in triplicates and their mean values are given.

RESULT AND DISCUSSION

Plant possess a well organized Reactive Oxygen Species (ROS) scavenging systems comprising enzymatic such as Catalase, Ascorbate peroxidase and Superoxide Dismutase, and non-enzymatic antioxidants. A coordinated function of these systems plays an important role in scavenging ROS and maintaining redox status of the cell. Relative to control seedlings, growth of mung seedlings was reduced due to Mn at tested concentrations.

Increased level of antioxidative enzymes protects the cell against the oxidative damage by removal of free radicals or reactive oxygen species. Catalase activity was calculated in control and stressed seeds at alternate days of germination. It was observed that the activity of catalase increased with increase in time as compared to unstressed seedlings. Increased level of catalase was observed, which showed that it is major enzyme in scavenging cellular H₂O₂

[11]. This enzyme is regarded as bioindicators of heavy metal toxicity and play important roles in scavenging ROS like H₂O₂ to reduce oxidative damage.


Ascorbate peroxidase activity was calculated in control and stressed seeds at alternate days of germination. It was observed that the activity increased with increase in time as compared to unstressed seedlings.

Activity of peroxidase increased with increase in time as compared to unstressed seedlings at alternate days of germination. The POX has been implicated in the synthesis of lignin and other phenolic polymers. So the enhancement in POX activity might defend the cells against harmful concentrations of hydroperoxides thereby protecting cellular components such as proteins and lipids against oxidation.


Lipid peroxidation occurs in plant tissues leading to production of free radicals, which are normally detoxified by antioxidative enzymes. Under stress conditions, more production of free radicals takes place. MDA is product of lipid peroxidation. So, lipid peroxidation can be estimated by measuring the level of MDA. In this study, it was observed that MDA content was high in germinating mung bean when treated with Mn as compared to control. So,

it was suggested that involvement of free radicals in membrane lipid peroxidation in the seeds subjected to Mn-stress could be a reason for increase in MDA content. Generally, free radical generation and membrane damage would be low in tolerant plants and thereby the formation of lower levels of MDA content.

SOD is a key enzyme in cell which plays a vital role against oxidative damage and severe environmental conditions. It was observed that at alternate days of germination the activity increased with increase in time as compared to unstressed seedlings.




(a)




(b)

It was observed that at alternate days of germination the Hydrogen peroxide content decreased with increase in time as compared to unstressed seedlings.


Protein content in Mn-treated seedlings was increased, indicating that heavy metal stress may induce production of stress proteins [12] including some heat shock proteins [13]. It may be suggested that protein content increased due to de novo synthesis of stress proteins provoked by metal exposure [14]. Several researchers found increased protein synthesis under metal stress.




(c)




(d)



(e)



(f)



(g)

Figure 1: Effect on antioxidant enzymes from germinated mung beans germinated under stress conditions (a) Catalase, (b) Ascorbate peroxidase, (c) Peroxidase, (d) MDA, (e) SOD, (f) Hydrogen peroxide, and (g) protein content

CONCLUSION

In conclusion, antioxidant machinery plays an important role in Mn-stress tolerance of plants. In the present study, we hypothesized that enhanced levels of antioxidants and MDA content on Mn exposure activate the multi tolerance mechanism of antioxidative enzymes under stress. The results of the present study also demonstrate the effect of Mn exposure, the cultivar capability to activate multi defense mechanism against oxidative damage caused by Mn ions may be a key factor in the detoxification mechanism of plant tolerance to unfavourable conditions.

ACKNOWLEDGEMENT

The authors are grateful to the management of SD College, Ambala Cantt for the infrastructural facilities provided to carry out the work.

REFERENCES

- [1] Malar, S., Vikram, S.S., Favas, P.J.C. and Perumal, V. (2014) Lead Heavy Metal Toxicity Induced Changes on Growth and Antioxidative Enzymes Level in Water Hyacinths [*Eichhornia crassipes* (Mart.)]. *Botanical Studies*, **55**, 1-11.
- [2] Puschenreiter, M., Schnepf, A., Molina-Millan, I., Fitz, W.J., Horak, O., Klepp, J., Schrefl, T., Lombi, E. and Wenzel, W.W. (2005) Changes

of Ni Biogeochemistry in the Rhizosphere of the Hyperaccumulator *Thlaspi goesingense*. *Plant and Soil*, **271**, 205-218.

- [3] Gupta V K, Sharma S K. 2006. Plants as natural antioxidants. *Natural Product Radiance*, Vol. 5(4), 326-334
- [4] Rani., P., U. K. Meena and J. Karthikeyan, 2004. Evaluation of antioxidant properties of berries. *Ind. J. of Clini. Biochem.*, 19(2):-103-110.
- [5] Aebi H (1984). Catalase in vitro. *Method Enzymol* 105: 121-126
- [6] Zhu Z, Wei G, Li J, Qian Q, Yu J (2004) Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.) *Plant Science* **167**: 527-533.
- [7] Shannon LM, Kay E, Lew JY (1966) Peroxidase isoenzymes from horseradish roots. *J. Biol. Chem.* 241: 2166-2172.
- [8] Sinha KA 1972 Colorimetric assay of catalase. *Anal Biochem* 47: 389-394.
- [9] Moshaty EFIB, Pike SM, Novacky AJ, Sehgal OP (1993) Lipid peroxidation and superoxide production in cowpea (*Vigna unguiculata*) leaves infected with

- tobacco ringspot virus or southern bean mosaic virus. *Physiol Mol Plant Pathol* **43**: 109-119.
- [10] Lowry OH, Rosenbrough HJ, Farr AL, Randall RJ (1951) Protein measurement with Folin-Phenol reagent. *Journal of Biological Chemistry* **193**: 265-275.
- [11] Vranova E, Inze D, Van Breusegem F (2002). Signal transduction during oxidative stress. *J Exp Bot* **53**: 1227–1236.
- [12] Sanita di Toppi L, Gabbrielli R (1999). Response to cadmium in higher plants. *Environ Exp Bot* **41**: 105–130.
- [13] Namjooyan S, Khavari-Nejad R, Bernard F, Namdjoyan S, Pir H (2012). The effect of cadmium on growth and antioxidant responses in the safflower (*Carthamus tinctorius* L.) callus. *Turk J Agric For* **36**: 145–152.
- [14] Verma S, Dubey RS (2003). Lead toxicity induces lipid peroxidation and alters the activities of antioxidant enzymes in growing rice plants. *Plant Sci* **164**: 645–655.