IJBPAS, December, 2026, 15(12): 6052-6072

ISSN: 2277–4998

International Journal of Biology, Pharmacy and Allied Sciences (IJBPAS)

'A Bridge Between Laboratory and Reader'

www.ijbpas.com

EXPLORING THE ROLE OF DARK CHOCOLATE IN REDUCING MENSTRUAL DISCOMFORT: COMPREHENSIVE REVIEW

BAIDYA M*, DAS A AND PAUL T

Milestones Institute of Pharmaceutical Sciences, Udaipur, Tripura-799120

***Corresponding Author: Dr. Moushumi Baidya: E Mail: baidyamoushhumi@gmail.com**

Received 29th May 2026; Revised 10th July 2026; Accepted 1st Oct. 2026; Available online 1st Dec. 2026

https://doi.org/10.31032/IJBPAS/2026/15.12.10689

ABSTRACT

Dysmenorrhea, a common problem that can disrupt a woman's daily activities and overall sense of well-being, is the pain associated with a woman's period. In recent times, dietary interventions like dark chocolate have gained popularity as a potential natural therapy for menstruation pain. Dark chocolate has magnesium, flavonoids, and other bioactive chemicals that may help ease symptoms like cramping, mood swings, and fatigue. In this article, we have reviewed the physiological processes by which dark chocolate may assist to decrease menstruation pain. Research has shown that regularly eating large amounts of high-cocoa dark chocolate can help to improve blood flow, reduce inflammation and relax muscles, all of which can help to decrease the amount of pain you feel. The aim of this review is to explore role of dark chocolate in alleviating menstrual discomfort, discuss the underlying biological mechanisms, summarize the current scientific evidence and discuss direct implications for its use.

Keywords: Dysmenorrhea, Menstrual cramps, Herbs, Ginger, Turmeric, Dark chocolate, Natural remedy, Anti-inflammatory, Women's health

INTRODUCTION

Dysmenorrhea, also referred to as menstrual cramps, is the term used to describe the discomfort that occurs during menstruation as a result of the uterus's muscular contractions. These contractions may put pressure on surrounding blood vessels, temporarily reducing the uterus's oxygen supply and causing pain and cramping **[1, 2]**.

The uterine lining's endogenous prostaglandins, which are often raised during menstruation, are the main source of this discomfort **[3]**. Menstrual cramps, a distressing sensation that many women go through every month, have been compared to the discomfort of a heart attack **[4]**.

Dysmenorrhea, or period pain, is common among women of reproductive age, including teenage girls, and can cause discomfort and interfere with everyday activities, demanding treatment **[5, 6]**. Chocolate includes copper, which produces neurotransmitters like as collagen and endorphins, acting as an analgesic and natural sedative to relieve pain **[7]**. The purpose of this study was to review the effect of dark chocolate on decreasing dysmenorrhea in adolescents.

A teenager is an individual with a mature anatomy and reproductive function. Adolescent puberty is distinguished by increasing levels of luteinizing hormone (LH) and follicle stimulating hormone (FSH), which cause the formation of sex hormones **[8]**. These hormones mature reproductive organs such as the ovaries, uterus, vagina, and breasts, as well as the formation of secondary sexual traits like as hair in the armpits, genitals, fine hairs on the skin, and face, among other things. This also marks the beginning of menstruation (menarche). Teenagers will experience discomfort before and throughout their menstrual periods, regardless of their cycle or timing. Menstrual pain, also known as dysmenorrhea, is stiffness or convulsions in the lower abdomen caused by menstruation and the generation of prostaglandins **[9]**. Pain usually begins the day before the menstrual cycle and lasts two days, or until menstruation ends. Menstrual discomfort prevalence ranges from 15.8 to 89.5%, with teenagers having the highest incidence **[10]**. Menstrual discomfort, which manifests as abdominal cramps, lower back pain, exhaustion, and mood swings, is mostly caused by increased production of prostaglandins, which cause uterine contractions. Nonsteroidal anti-inflammatory medicines (NSAIDs), heat therapy, and lifestyle changes are examples of conventional treatments. Concerns about long-term pharmaceutical use have driven interest in natural food therapies **[11]**.

Dysmenorrhea, is one of the most frequent gynaecological issues among adolescents and women of reproductive age. It is distinguished by intense menstrual cramps that occur before or during menstruation and can be accompanied by symptoms such as nausea, fatigue, headache, dizziness, and lower back discomfort. According to studies, dysmenorrhea affects 45% to 95% of menstrual women worldwide, making it a major public health concern **[12]**. The prevalence varies according to age, lifestyle, cultural variables, and diagnostic criteria.

Severe menstruation discomfort can interfere with daily activities, academic performance, work productivity, social contacts, and physical activity **[13]**. Many people report being absent from school or work during menstruation, while others have difficulty concentrating and working efficiently even when they are present **[14]**. In addition to physical limits, dysmenorrhea is linked to psychological symptoms such as impatience, worry, tension, and mood swings, which can further impair general well-being.

According to research, recurring menstruation pain can have a negative impact on sleep quality, emotional health, and interpersonal connections. The burden of dysmenorrhea may result in lower life satisfaction and health-related quality of life, especially among those who experience moderate to severe symptoms **[14]**. Given its high prevalence and pervasive implications, effective management measures are required to lessen the physical, mental, and socioeconomic consequences of menstruation discomfort and improve the overall quality of life for those affected **[15]**. Menstrual discomfort is a frequent health issue that impacts the daily lives and well-being of millions of women worldwide. Despite its widespread presence, many people seek simple, cost-effective, and natural ways to treat menstruation pain **[16]**. Dark chocolate has received attention as a potential dietary cure due to its high concentration of magnesium, flavonoids, and other bioactive chemicals that may help relieve cramps and enhance mood. Exploring the scientific evidence supporting this widely available meal allows us to better appreciate its possible function in menstrual health. This topic includes nutrition, women's health, and preventative care, making it important to contemporary research and public health awareness **[17]**.

Conventional therapies for menstrual pain generally include nonsteroidal anti-inflammatory drugs (NSAIDs) or hormonal treatments, but there is increasing interest in natural and nutritional approaches as adjunct therapies **[18]**.

Dark chocolate is a good source of magnesium, flavonoids and other bioactive compounds that have become of interest as a possible treatment for dysmenorrhea. Magnesium may help relax uterine muscles and lower the severity of menstrual cramps **[19]**. Antioxidants included in dark chocolate can help decrease inflammation and improve blood flow. Dark chocolate also releases endorphins and serotonin which can increase mood and lessen menstruation discomfort. Dark chocolate as a generally accepted and readily available food product is a potential natural alternative to enhance the dysmenorrhea treatment and to improve the menstrual well-being **[20]**.

The purpose of this study is to explore the possibility that dark chocolate may ease menstrual discomfort by analyzing the current evidence on its bioactive constituents, mechanisms of action and efficacy for the treatment of dysmenorrhea. The goal is to investigate whether dark chocolate could be a good complement for menstrual pain therapy **[21]**.

Many women experience period pain before and throughout their period. This is medically referred to as dysmenorrhea. The symptoms can vary widely in severity, duration and impact and some people will find them more disruptive to their normal life than others **[22]**.

1.1 Types of Dysmenorrhea

• **Primary Dysmenorrhea**: Primary dysmenorrhea is menstruation pain in females with normal pelvic structure, generally starting in adolescence. It is said that this condition occurs only in ovulatory cycles and usually appears between 6 and 12 months after the commencement of menstruation and without any known pathology or organic etiology **[28]**. The exact etiology of primary dysmenorrhea is unknown, but it is connected with increased prostaglandin levels during the ovulatory cycle.

• **Secondary Dysmenorrhea**: Secondary dysmenorrhea is menstrual pain that occurs in association with underlying medical conditions such as endometriosis, adenomyosis, fibroids (myomas), and various other pelvic disorders **[29]**.

1.2 Causes of Dysmenorrhea:

- **Hormonal fluctuations:** Levels of estrogen and progesterone change during the menstrual cycle. These hormones drop just before menstruation and this triggers the uterine lining to shed. Most of the symptoms including period cramps are because of the drop in progesterone **[23]**.
- **Release of prostaglandins:** Prostaglandins are hormone-like molecules that the uterus releases to help push out the uterine lining, and to create muscular contractions. The prostaglandins go up, you experience headaches, nausea, diarrhea, severe cramps **[24]**.
- **Underlying health issues:**

  Endometriosis: Tissue that normally lines the inside of your uterus grows outside your uterus. brings on horrible discomfort, and even fertility issues.

  Uterine fibroids: Uterine fibroids are benign growths in the uterus that may cause pain, cramping and bleeding **[25]**.

  Pelvic inflammatory disease (PID): It is an infection of the reproductive organs that may worsen symptoms of menstruation and induce pain.

Adenomyosis: The tissue that normally lines the uterus grows into the muscle of the uterus, leading to severe cramping. The illness causes agonizing cramps **[26]**.

- **Genetic and Lifestyle Factors:**

Cramps can be hereditary – that is, it runs in your family.

Early menarche (before 11 years), copious bleeding and prolonged periods are associated with more severe symptoms **[27]**.

Lack of exercise, stress and smoking might worsen the problem.

1.3 Symptoms common among women

Physical Signs:

- Abdominal cramps are painful and can be felt in the lower abdomen and vary in strength. They could come to the back of the thighs.
- Low Back Pain.
- Hormones can fluctuate and cause migraines and headaches.
- Excess prostaglandin makes you nauseous and vomit.
- Bloating is a condition that causes fluid retention, leading to a feeling of heaviness or distention in the belly.
- Tenderness in the breast: Swelling or pain before or during your period.
- Problems with digestion. You can get constipation or diarrhea.
- Unusual fatigue or weakness.
- Sometimes dizziness occurs because of low iron or blood loss **[28-30]**

Behavioural and Emotional Cues:

- Mood Swings: Mood swings, impatience, or sensitivity.
- Gloom or anxiety Some people are more emotionally distressed during their menstruation **[31]**.
- Sleep problems: Difficulty in falling asleep due to pain or discomfort.
- Appetite Changes: Cravings or a reduced appetite might be experienced during menstruation.
- Menstrual pain and effect on daily activities.
- Achievements (Professional and Academic).

Social & Family Life:

- Pain and sorrow may induce one to avoid social events or family gatherings.
- Communication and relationships can be affected by emotional challenges **[32]**.
- Mental Wellness: The distress you feel right now can make ongoing stress, worry or low self-esteem worse.
- They can also happen with illnesses like premenstrual dysphoric disorder (PMDD), a condition that causes significant mood swings **[33]**

1.4 Mechanisms of Dysmenorrhoea

Several physiological processes that take place during menstruation are the cause of menstrual discomfort or primary dysmenorrhea: Increased prostaglandin synthesis, excessive uterine contractions, and inflammation with oxidative damage are the primary mechanisms. The typical menstrual cramping ache is produced when they unite **[34]**.

Prostaglandin production

Prostaglandins are hormone-like substances produced by the endometrium (lining of the uterus). During menstruation a lot of prostaglandins are released due to destruction of endometrial cells **[35]**. Mostly prostaglandin F2α (PGF2α) and prostaglandin E2 (PGE2). These chemicals in high doses cause strong contractions of the uterine muscle and narrowing of the blood vessels of the uterus. Those with severe menstrual pain have been found to have a higher level of prostaglandins than those with mild or no symptoms. Thus, it is believed that increased prostaglandin production is one of the major causes of dysmenorrhea **[36]**.

Uterine contraction

This causes the uterine muscles to contract more often and more powerfully. These contractions help to push out the lining of the uterus during your period. But too much contraction can squeeze the arteries around it and limit blood flow to the uterine tissue **[37]**. This means that the delivery of oxygen to the uterus (uterine ischaemia) is less. This makes the cramping agony worse. The stronger the cramps during the period, the stronger and longer the uterus contracts.

Oxidative stress and inflammation

Inflammatory mechanisms are important in dysmenorrhea. During menstruation, cytokines, prostaglandins and other inflammatory mediators are released to allow local inflammation in the uterus. This inflammation can cause you to be more sensitive to pain and cause symptoms such as stomach cramping, headache, fatigue and nausea **[38]**. Oxidative stress is defined as an imbalance between the production of reactive oxygen species (free radicals) and antioxidant defenses of the body. Research suggests that people with dysmenorrhea may experience increased oxidative stress, which can contribute to more inflammation and tissue injury. Increased oxidative stress may also increase pain perception and contribute to the severity of menstrual symptoms **[39]**. So, food rich in antioxidants and minerals have been studied as a possible treatment in the diet to ease menstruation discomfort. The main molecular players of menstruation pain are overproduction of prostaglandins, strong uterine contractions and oxidative stress caused by inflammation. Understanding these mechanisms is important for the development of adequate treatment and nutritional strategies in the management of dysmenorrhea **[40]**.

Psychosocial and Emotional Manifestations

In addition to physical discomfort, dysmenorrhea can also produce significant psychosocial and emotional consequences. Menstrual pain is recurrent and may affect emotional well-being, daily functioning and general quality of life. Many people experience mood problems, increased stress and symptoms of anxiety throughout their menstrual cycle, often with dysmenorrhea **[41]**.

Mood Changes

Changes in mood is one of the most common emotional signs of dysmenorrhea. The hormonal changes that occur during the menstrual cycle along with the physical pain and discomfort can make you irritable, unhappy, moody and emotionally sensitive. The pain of menstruation can also lead to reduced social, academic and work-related activities, which can cause frustration and reduced life satisfaction. These emotional responses may help to contribute to the overall experience of pain and establish a vicious cycle of pain and mood disorders exacerbating one another **[42]**.

Anxiety and Stress Disorder

Such patients are more prone to high levels of stress and anxiety. The anticipation of repeated menstrual pain can cause psychological distress even before the onset of menstruation. Women with mild to severe dysmenorrhea may worry about their ability to perform daily tasks, attend school or work or engage in social activities during their menstrual period. This worry can be stressful and may contribute to symptoms of anxiety. Research has also linked chronic pain with activation of the body's stress response system, which can increase pain sensitivity and damage emotional health **[43]**. Psychological factors such as stress, anxiety and negative emotions have been proven to have an impact on pain perception and aggravate the severity of menstruation symptoms. The physical and the mental components of the dysmenorrhea disorder should therefore be considered in the effective treatment of dysmenorrhea. Approaches that reduce pain, improve mood, and promote relaxation may be important to enhance overall menstrual health and quality of life **[44]**.

1.5 Physiology of menstrual cycle

Menstruation is a biological process. It involves a series of changes controlled by female hormones, ending in periodic bleeding. Some women may have menstruation problems, such as heavy bleeding. The menstrual cycle can be divided into four main phases: 1) Menstruation (Day 1 to Day 5) 2) Follicular phase (Day 1-13) 3) Ovulation (Day 14) 4) Luteal phase (Day 15-28).




Figure 1: Menstrual cycle

2. Role of Dark Chocolate

Dark chocolate is created with less milk tha n milk chocolate and a larger percentage of cocoa butter and solids.

On the packaging, the proportion of cocoa c ontent is often indicated. Dark chocolate's nutritional and bioactive components may help lessen dysmenorrhea.

Classification of dark chocolate

A number of criteria, including as cocoa content, ingredients, and quality, may be used to categorize dark chocolate. This is a summary of the common classifications for it:

1. By Content of Cocoa: The color and bitterness of the chocolate are determined by the proportion of cocoa solids (cocoa mass + cocoa butter).

2. By Composition of Ingredients: Unlike milk chocolate, all dark chocolate is free of milk solids; nonetheless, there are differences depending on the ingredients that are added.

• Pure Dark Chocolate: just sugar, cocoa butter, and cocoa mass.

Dark chocolate that has been flavored with nuts, fruit (orange, raspberry), or spices (cinnamon, chili).

• Single-Origin Dark Chocolate: Usually featuring distinctive taste qualities, this kind of chocolate is made from cocoa beans from a single area or farm **[45]**.

3. By Production Method/Quality

• Artisan/Craft Chocolate: Bean-to-bar chocolate made in small batches with few ingredients and responsibly sourced beans.

• Commercial/Mass-Produced Chocolate: Produced in greater numbers, it may include additional sugar and emulsifiers such soy lecithin.

• Raw Dark Chocolate: Made from unroasted cacao beans, this chocolate has a unique flavor and retains more nutrients and antioxidants.

4. By Industry or Legal Standards (depending on the nation) Certain areas use a minimum cocoa content to describe dark chocolate:

• EU: Cocoa solids must be at least 35%.

• USA (FDA): There is no minimum, however it cannot be classified as "dark" or "semisweet" if it contains milk solids.

Dark Chocolate's Nutritional Profile

Dark chocolate is a great source of vital nutrients and bioactive chemicals since it is made from cocoa beans and has a higher proportion of cocoa solids than milk chocolate. Dark chocolate's nutritional makeup varies according to its cocoa level, however goods with 70% or more cocoa are often thought to be the healthiest **[46]**.

1. **Magnesium**

Magnesium, a mineral involved in over 300 enzymatic activities in the body, is abundant in dark chocolate. Magnesium is essential for nerve function, muscular relaxation, and the control of inflammatory reactions. Magnesium is especially important for period health since it may help lessen uterine muscle spasms and menstrual cramps **[47]**.

2. **Polyphenols and Flavonoids**

Flavonoids, particularly catechins, epicatechins, and procyanidins, are abundant in dark chocolate. These substances have potent anti-inflammatory and antioxidant qualities that aid in shielding cells from oxidative damage. By increasing nitric oxide production and blood flow, flavonoids also promote vascular health, which may lessen menstruation discomfort and enhance circulation.

3. **Iron**

The synthesis of haemoglobin and the movement of oxygen throughout the body depend on iron, an important mineral. Dark chocolate's iron content may assist boost daily iron intake and lessen period tiredness since menstrual blood loss might raise the risk of iron insufficiency **[48]**.

4. **Other Minerals and Potassium**

Dark chocolate contains iron, magnesium, potassium, copper, phosphorus, zinc, and manganese. These minerals support a number of physiological processes, such as immunological response, energy metabolism, muscular contraction, and antioxidant defenses systems **[49]**.

5. Caffeine and theobromine

Theobromine and trace levels of caffeine are naturally present in dark chocolate. As a moderate stimulant, theobromine may enhance mood, alertness, and general well-being. These substances may improve energy levels and lessen exhaustion during menstruation, although excessive use may make some people sensitive **[50]**.

5. Good Fats

Monounsaturated and saturated fats, namely oleic acid, stearic acid, and palmitic acid, are found in the cocoa butter found in dark chocolate. While oleic acid is linked to cardiovascular advantages, stearic acid is thought to have no effect on blood cholesterol levels **[51]**.

6. Fiber in the Diet

Dietary fiber, which is found in dark chocolate, aids with blood sugar regulation, satiety, and digestive health. Compared to milk chocolate variants, chocolates with a higher cocoa content often have more fiber **[52]**.

7. Antioxidants

One of the best dietary sources of antioxidants is dark chocolate. These substances may help decrease inflammatory levels, lessen oxidative stress, and neutralize free radicals. One of dark chocolate's main health benefits is thought to be its antioxidant activity **[53]**.

Significance for Menstrual Health

Dark chocolate is a potentially helpful food during menstruation since it contains iron, flavonoids, magnesium, and other bioactive substances. These nutrients may aid in uterine muscle relaxation, inflammation reduction, mood enhancement, tiredness prevention, and general wellbeing. Dark chocolate's nutritional composition indicates that moderate intake may be a helpful dietary approach for reducing menstruation pain, even if it shouldn't be used as a substitute for medical therapy **[54]**.

3. Mechanisms Through Which Dark Chocolate May Reduce Menstrual Discomfort

Magnesium, flavonoids, polyphenols, and antioxidants are among the bioactive substances found in dark chocolate that may reduce menstruation pain via a number of physiological processes. These elements may lessen the intensity of dysmenorrhea by affecting blood circulation, inflammation, muscle function, and pain pathways **[55]**.

3.1 Uterine Muscle Relaxation

Promoting uterine muscle relaxation is one of the main ways that dark chocolate may lessen menstruation discomfort. Magnesium, a vital element involved in nerve and muscle function, is abundant in dark chocolate. Magnesium aids in controlling the flow of calcium into muscle

cells. Magnesium functions as a natural muscle relaxant by reducing excessive calcium activity since calcium causes muscular spasms. This impact may lessen the frequency and force of uterine contractions during menstruation, which may lessen the discomfort associated with cramps. Increased menstrual comfort and less dysmenorrhea symptoms have been linked to adequate magnesium consumption **[56]**.

3.2 Lowering of Inflammatory Reactions

Menstrual discomfort is mostly caused by inflammation. Inflammatory mediators are generated in the uterus during menstruation, which exacerbates tissue irritation and discomfort. Flavonoids and polyphenols found in dark chocolate have potent anti-inflammatory qualities. These drugs lessen the synthesis of chemicals that cause inflammation and aid in the suppression of inflammatory pathways. Dark chocolate may help reduce the intensity of menstrual cramps and other related symptoms including exhaustion and discomfort by reducing inflammatory activity **[57]**.

3.3 Prostaglandin Activity Modulation

During menstruation, prostaglandins, which similar to hormones, cause uterine contractions. One of the main causes of primary dysmenorrhea is excessive prostaglandin production, which results in greater contractions, decreased blood flow, and higher discomfort. Dark chocolate has anti-inflammatory and antioxidant substances that may help control prostaglandin production and activity. Dark chocolate's bioactive ingredients may help lower excessive prostaglandin-mediated reactions, which might lessen period pain and discomfort even if it doesn't totally stop prostaglandin synthesis **[58]**.

3.4 Improvement of Vascular Function and Blood Flow

Tissue ischemia and greater discomfort may arise from reduced blood supply to the uterus during strong contractions. Cocoa flavonoids, which are abundant in dark chocolate, are known to enhance vascular function by increasing nitric oxide generation. Vasodilation is facilitated by nitric oxide, which causes blood vessels to relax and enlarge. In addition to improving oxygen supply to tissues, improved circulation may lessen menstrual ischemia discomfort. By lessening the impact of limited uterine circulation, improved blood flow might help promote general cardiovascular health and lessen monthly pain **[59]**.

When taken as a whole, these pathways imply that dark chocolate might be a helpful dietary choice for controlling menstruation pain. A biological foundation for its possible advantages in lowering menstruation pain is provided by its magnesium concentration, anti-inflammatory qualities, effects on

prostaglandin activity, and capacity to increase blood circulation **[60]**.

3.5. Modulation of Mood and Pain Perception

By boosting the production of endorphins and serotonin, two neurotransmitters linked to better mood and pain alleviation, dark chocolate may affect the central nervous system. During menstruation, these neurochemical alterations might lessen pain perception and enhance mental health. Theobromine's modest stimulant effects could also help with increased alertness and decreased weariness **[61]**.

3.6. Antioxidant Defenses Against Oxidative Damage

Increased inflammation and pain sensitivity are linked to oxidative stress. Antioxidants, especially flavonoids, which counteract free radicals and lessen oxidative damage, are abundant in dark chocolate. During menstruation, this antioxidant activity may help reduce inflammatory reactions and promote general uterine health **[62]**.

4. Proof from Scientific Research

In recent years, there has been an increase in scientific interest in the connection between menstruation pain and dark chocolate. The minerals included in dark chocolate may help lessen menstruation discomfort, according to research on magnesium supplements, clinical trials, and observational studies. This article reviews the available data and assesses its benefits and drawbacks **[63]**.

4.1 Observation studies

The relationship between dietary practices and menstruation symptoms in adolescents and women of reproductive age has been investigated via observational studies. According to a number of studies, those who eat a diet high in antioxidants, magnesium, and anti-inflammatory foods often have less severe menstruation discomfort. Researchers have proposed a possible connection between eating dark chocolate and better menstruation comfort since it contains these nutrients **[64]**.

According to some studies, women often eat chocolate during their periods in order to control pain, enhance their mood, and satisfy cravings. Although these results point to a potential positive link, observational studies are unable to prove a clear cause-and-effect relationship due to the involvement of individual, dietary, and lifestyle variables.

4.2 Clinical Research on Menstrual Pain and Dark Chocolate

The effects of dark chocolate on dysmenorrhea have only been the subject of a small number of clinical research. According to the results of this research, eating dark chocolate with a high cocoa content may lessen the severity of menstruation discomfort. When compared to baseline readings, those who ate dark

chocolate during their periods often reported reduced pain levels and increased comfort.

The magnesium and flavonoid content of dark chocolate, which may help ease uterine muscles, lessen inflammation, and enhance blood circulation, are the main causes of these benefits, according to researchers. Dark chocolate may provide both psychological and physical advantages during menstruation, according to certain research that have also shown increases in mood and emotional well-being.

Nevertheless, the majority of clinical studies have had rather small sample sizes and brief intervention durations, which restricts how broadly their results may be applied.

4.3 Data from Research on Magnesium Supplementation

Numerous studies have looked at how supplementing with magnesium affects menstruation pain. By controlling calcium activity in muscle cells, magnesium has been shown to lessen the frequency and intensity of uterine contractions. Supplementing with magnesium has been shown to enhance overall menstrual symptoms, decrease prostaglandin production, and lessen cramping.

These results suggest that dark chocolate may be useful in treating dysmenorrhea since it is naturally high in magnesium. Even while dark chocolate typically contains less magnesium than supplements, frequent consumption may increase magnesium intake and menstrual health.

4.4 Comparative Evaluation of Results

Dark chocolate may be beneficial for menstruation pain, according to the results of observational studies, clinical trials, and magnesium research taken together. The most concrete evidence comes from clinical research, which show that eating dark chocolate lowers self-reported pain levels. Nutrient-rich diets are linked to less severe dysmenorrhea, according to observational studies, and magnesium supplementation provides a compelling scientific basis for these advantages.

Even with these positive results, the evidence is still not as strong as it is for traditional therapies like nonsteroidal anti-inflammatory medications (NSAIDs). It is challenging to get firm conclusions on the efficacy of dark chocolate as a therapeutic intervention due to variations in research design, cocoa concentration, serving sizes, and participant characteristics **[65]**.

4.5

Benefits and Drawbacks of Current Rese arch

• Offers first data in favor of a dietary strategy for managing menstruation discomfort.

• Shows biological plausibility based on the actions of flavonoids and magnesium.

• Offers further advantages for general wellbeing, stress relief, and mood.

• Draws attention to a nutritional alternative that is widely accessible, reasonably priced, and typically safe.

Drawbacks

• There aren't many studies that particularly address dark chocolate with dysmenorrhea.

• The statistical power of many studies is diminished by small sample sizes.

• Variability in chocolate intake duration, dose, and cocoa content between research.

• The use of self-reported pain measurements, which might be biased.

• Insufficient long-term research assessing safety and long-term consequences.

• It might be challenging to distinguish the effects of dark chocolate from other dietary and lifestyle variables.

According to recent research, dark chocolate's magnesium content and anti-inflammatory qualities may help lessen menstruation pain. However, more extensive, well planned randomized controlled studies are required to verify its efficacy, identify ideal consumption amounts, and provide precise guidelines for clinical practice.

5. Extra Advantages of Dark Chocolate for Menstruation

Dark chocolate may have a number of other advantages during menstruation in addition to its possible ability to lessen menstrual discomfort. Its abundance of bioactive substances, including flavonoids, magnesium, theobromine, and antioxidants, which may improve mood, energy, and general physiological well-being, is primarily responsible for these advantages.

5.1 Enhancement of Mood

Emotional swings, such as anger, melancholy, and mood swings, are often linked to menstruation. There are many ways that dark chocolate may elevate mood. It increases the production of endorphins, which are naturally occurring "feel-good" chemicals that lessen pain perception and increase sensations of pleasure. Furthermore, serotonin, a neurotransmitter strongly associated with mood regulation and emotional stability, may become more active in the brain after consuming dark chocolate.

Chocolate's delicious flavor and sensory appeal may also provide psychological comfort by lowering emotional stress during menstruation. Many people's innate need for chocolate throughout their menstrual cycle may be explained by these combined effects.

5.2 Fatigue Reduction

During menstruation, fatigue is a typical symptom that is often brought on by hormonal changes, blood loss, and low energy. Small levels of theobromine and caffeine, two moderate stimulants that might improve alertness and lessen fatigue, can be found in dark chocolate. These substances may increase physical vigor and mental clarity without having significant stimulant effects.

Dark chocolate's iron concentration may also aid in the body's oxygen delivery, especially for those with low iron levels brought on by menstrual blood loss. This may also help to enhance vigor and lessen exhaustion during menstruation.

5.3 Cardiovascular and Antioxidant Advantages

One of the best food sources of antioxidants, especially flavonoids like catechin and epicatechin, is dark chocolate. These substances aid in the body's reduction of oxidative stress and neutralization of free radicals. Reducing oxidative stress may indirectly enhance menstruation comfort since it is linked to inflammation and pain sensitivity.

By enhancing endothelial function and encouraging the synthesis of nitric oxide, which helps relax blood vessels and enhance circulation, flavonoids in dark chocolate also promote cardiovascular health. Improved blood flow may lessen the ischemic discomfort brought on by menstrual cramps and promote general uterine health.

5.4 Possibility of Enhancing General Well-Being

An overall feeling of well-being during menstruation is influenced by the combined effects of mood improvement, decreased tiredness, antioxidant protection, and enhanced circulation. During their menstrual cycle, dark chocolate may make people feel more at ease, at ease, and emotionally balanced.

These advantages emphasize the potential importance of dark chocolate as a supplemental dietary choice for enhancing both the physical and emotional elements of menstrual health, even though they are supportive rather than curative. It may improve general quality of life during menstruation when used in moderation.

6. Suggested Cocoa Amount

The proportion of cocoa in dark chocolate has a direct bearing on its possible advantages. Generally speaking, products with 70% or more cocoa are advised since they provide higher concentrations of flavonoids, magnesium, and other beneficial substances. Chocolates with a lower cocoa content often have less beneficial phytochemicals and more added sugar, which may lessen their health benefits.

6.2 Recommended Intake Amounts

When eating dark chocolate to relieve menstruation pain, moderation is crucial. Depending on dietary requirements and individual tolerance, a typical recommended daily consumption is between 20 and 40 grams. This quantity is thought to be enough to maximize possible advantages while reducing excessive consumption of calories and sugar. Because of its high energy density, overconsumption should be avoided.

During the premenstrual period and the first few days of menstruation, when symptoms like cramps, exhaustion, and mood swings are at their worst, dark chocolate may be especially helpful. To maintain constant magnesium levels and antioxidant intake, some people may also benefit from frequent moderate ingestion during the menstrual cycle.

6.3 Combining Other Lifestyle Interventions with Dark Chocolate

Dark chocolate should not be used as a stand-alone therapy, but rather as a supplementary strategy. When paired with other healthful lifestyle choices like these, its advantages can be amplified.

• Frequent exercise (e.g., stretching)
• Sufficient hydration
• A well-balanced diet full of healthy grains, fruits, and veggies
• Stress-reduction methods like yoga or meditation
• Good sleep hygiene

7. Prospects for Further Research

Future research should concentrate on the following to have a deeper understanding of dark chocolate's impact on menstrual health:

• Extensive randomized controlled studies to provide more robust clinical data
• The ideal dose and time to ascertain the levels of effective intake
• Mechanistic research to elucidate the manner in which cocoa chemicals impact inflammatory and hormonal processes
• Research comparing the relative efficacy of traditional treatments like NSAIDs and magnesium supplements

Clearer nutritional guidelines and therapeutic applications will result from such study.

CONCLUSION

Menstrual discomfort is a common condition that may negatively affect a person's physical, emotional, and social well-being. Even though traditional remedies like heat therapy, exercise, and nutritional supplements are still important options, growing evidence suggests that dark chocolate may be a helpful additional method. The bioactive components of dark chocolate, including as magnesium, iron, and flavonoids, may help reduce menstrual pain by increasing circulation, lowering inflammation, relaxing muscles, and stabilizing mood. Consuming dark chocolate, particularly those with 70% or more cacao, has been demonstrated in clinical studies to reduce the intensity of menstrual pain. Compared to many alternative medicines, dark chocolate has many advantages, such as being easily available, affordable, easy to consume, and generally having minimal side effects when taken in moderation.

Even while its pain-relieving effects may not be as potent as those of prescription medications like NSAIDs, it may significantly alleviate mild to moderate

menstrual discomfort and improve mental well-being during menstruation.

In conclusion, dark chocolate is a useful, affordable, and pleasurable dietary approach that may lessen menstruation pain. Further studies will provide additional insight on its efficacy, ideal dose, and function in the overall management of menstrual health.

REFERENCES

[1] Grandi G., Ferrari S., Prevalence of menstrual pain in young women: what is dysmenorrhea, Journal of Pain Research, 2012, 5, 169–174.

[2] De Souza A., Mendieta M., Menstrual cramps: A new therapeutic alternative care through medicinal plants, Health, 2013,5(7),1106 1109.

[3] Azizah R., Anggreini P., Effect of Combination of Dark Chocolate and Herbal Ingredients for dysmenorrhea in Late Adolescents, Journal of Tropical Pharmacy and Chemistry, 2023, 7(1).

[4] Verma A., Kadam A., Does Dark Chocolate Relieve Menstrual Pain in Adult Women?: A Study Among Indian Population, International Journal of Physiology, 2019, 7(4), 16-21.

[5] Iacovides, S., Avidon, I., & Baker, F. C. (2015). What we know about primary q bnm

[6] Itani, R., Soubra, L., Karout, S., Rahme, D., Karout, L., & Khojah, H. M. J. (2022). Primary Dysmenorrhea: Pathophysiology, Diagnosis, and Treatment Updates. *Korean Journal of Family Medicine*, *43*(2), 101-108.

[7] Martire, F. G., Piccione, E., Exacoustos, C., & Zupi, E. (2023). Endometriosis and Adolescence:The Impact of Dysmenorrhea. *Journal of Clinical Medicine*, *12*(17), 5624.

[8] Santrock, J. W., & Santrock, J. W. (2007). *Psikologipendidikan [Educational psychology]* (2nd ed.) Jakarta: Kencana Prenada Media Group.

[9] Ergonul, P. G., Ergonul, B., & Seckin, A. K. (2010). Cholesterol content and fatty acid profile of chocolates mostly consumed in Turkey. *CyTA - Journal of Food*, *8*(1), 73–78.

[10] Razola-Díaz, M. C., Aznar-Ramos, M. J., Verardo, V., Melgar-Locatelli, S., Castilla-Ortega, E., & Rodríguez-Pérez, C. (2023). Exploring the Nutritional Composition and Bioactive Compounds in Different Cocoa Powders. *Antioxidants*, *12*(3), 716.

[11] Jalil, A. M. M., & Ismail, A. (2008). Polyphenols in Cocoa and Cocoa Products: Is There a Link between Antioxidant Properties and Health? *Molecules*, *13*(9), 2190–2219.

[12] Melo, T. S., Pires, T. C., Engelmann, J. V. P., Monteiro, A. L. O., Maciel, L. F., & Bispo, E. S. (2020). Evaluation of the content of bioactive compounds in cocoa beans during the fermentation process. *Journal of*

Food Science and Technology, *58*(5), 1947–1957.

[13] Quek, R. Y. C., Peh, E. W. Y., & Henry, C. J. (2020). Effects of Cocoa Butter and Cocoa Butter Equivalent in a Chocolate Confectionery on Human Blood Triglycerides, Glucose and Insulin. *Foods*, *9*(4), 455.

[14] Singh, P. K., Khedkar, R. D., & Chandra, S. (2024). Chocolate: An overview of functional potential and recent trends in fortification. *Brazilian Journal of Food Technology*, *27*.

[15] Soares, T. F., & Oliveira, M. B. P. P. (2022). Cocoa By-Products: Characterization of Bioactive Compounds and Beneficial Health Effects. *Molecules*, *27*(5), 1625.

[16] Kababie-Ameo, R., Rabadan-Chavez, G. M., Vazquez-Manjarrez, N., & Gutierrez-Salmean, G. (2022). Potential applications of cocoa (Theobroma cacao) on diabetic neuropathy: mini-review. *Frontiers in Bioscience-Landmark*, *27*(2).

[17] Bali, S. (2025). Zesty Pomegranate Bon-Bon Candies: A Tasty Boost For Your Brain And Gut Health. *International Journal of Medical and Pharmaceutical Research*, *6*(6), 461–475.

[18] Maharani, *et al*. Effect of dark chocolate consumption on menstrual pain among late adolescents. 2017. Volume 3, Issue 6. 686–692.

[19] Karakus Selcuk, Baysal, *et al*. Effect of dark chocolate combined with relaxation therapy on dysmenorrhea pain and anxiety. 2022. Volume 56 102192

[20] Ferina, *et al*. Impact of dark chocolate on dysmenorrhea among adolescents. 2023. Volume 11, Issue 2. 192–195.

[21] Fatkurrohman, *et al*. Effectiveness of dark chocolate in reducing dysmenorrhea in school-based adolescents. 2023. Volume 3, Issue 2. 105–111

[22] Okta Viantri, *et al*. Role of dark chocolate in managing dysmenorrhea among adolescents in educational settings. 2025. Volume 17, Issue 1. 48–54

[23] Firnandani S, Achyar K, *et al*. Dark chocolate consumption and its effect on dysmenorrhea: a systematic review. 2025. Volume 13, Issue 1. 47–58

[24] Maharani, *et al*. Effect of dark chocolate consumption on menstrual pain among late adolescents. 2017. Volume 3, Issue 6. 686–692.

[25] Karakus Selcuk, Baysal, *et al*. Effect of dark chocolate combined with relaxation therapy on dysmenorrhea pain and anxiety. 2022. Volume 56 102192

[26] Ferina, *et al*. Impact of dark chocolate on dysmenorrhea among adolescents. 2023. Volume 11, Issue 2. 192–195.

[27] Fatkurrohman, *et al*. Effectiveness of dark chocolate in reducing

dysmenorrhea in school- based adolescents. 2023. Volume 3, Issue 2. 105–111.

[28] Okta Viantri, *et al*. Role of dark chocolate in managing dysmenorrhea among adolescents in educational settings. 2025. Volume 17, Issue 1. 48–54.

[29] Firnandani S, Achyar K, *et al*. Dark chocolate consumption and its effect on dysmenorrhea: a systematic review. 2025. Volume 13, Issue 1. 47–58.

[30] Satriawati1 A., Nugraheny E., The Effect of Combination of Warm Compression and Chocolate Against Menstrual Pain Reduction (Dysmenorrhea) In Teens In SMP Negeri Bangkalan, Indonesian Journal of Nursing and Midwifery, 2020, 8(1), 36-42.

[31] Saidah1 S., Lathifah M., The Effect of Giving Dark Chocolate on Reducing Menstrual Pain in Primary dysmenorrhoea in Employees of Cabangbungin General Hospital, Bekasi regency in 2023, Bioscientia Medicina: Journal of Biomedicine & Translational Research, 2023, 3(2), 3260-3263.

[32] Arfailasufandi, R., & Andiarna, F. (2018). Pengaruh Pemberian Coklat Hitam terhadap Penurunan Nyeri Haid pada Dismenorhea Primer The Influence of Dark Chocolate to Reduce Menstrual Pain in Primary Dysmenorhea. Journal of Health Science and Prevention, 27 35.

[33] Maharani, S. I., Pramono, N., Wahyuni, S. (2017). Dark chocolate's effect on menstrual pain in late adolescents. Belitung Nursing Journal, 3(6), 686-692. https://doi.org/10.33546/bnj.292.

[34] Wulandari, S., & Afriliana, F. D. (2017). The effect of consumption dark chocolate against primary dismenore pain for girls teenager atKkediri 5 high School. Indian Journal of Medical Research and Pharmaceutical Sciences, 4(2), 17-24.

[35] Hajati, A., Brondani, M., Angerstig, L., Klein, V., Liljeblad, L., Al-Moraissi, E. A., Louca Jounger, S., Brondani, B., & Christidis, N. (2023). Chocolate intake and muscle pain sensation: A randomized experimental study. *PLOS ONE*, *18*(5), e0284769.

[36] Nuha, K., Rusmil, K., Ganiem, A. R., Permadi, W., & Diah Herawati, D. M. (2023). Single-Blind Randomized Controlled Trial: Comparative Efficacy of Dark Chocolate, Coconut Water, and Ibuprofen in Managing Primary Dysmenorrhea. *International Journal of Environmental Research and Public Health*, *20*(16), 6619.

[37] Nurazizah, E., Tih, F., & Suwindere, W. (2015). Black Chocolate Consumption Reduces Subjective Symptoms in 18-22 Years Old Females with Premenstrual

Syndrome. *Journal of Medicine & Health*, *1*(1).

[38] Safari, K., Hemmatinafar, M., Suzuki, K., & Jahromi, M. K. (2025). Dark Chocolate Mitigates Premenstrual Performance Impairments and Muscle Soreness in Female CrossFit® Athletes: Evidence from a Menstrual-Phase-Specific Trial. *Nutrients*, *17*(8), 1374.

[39] Sultana, A., Rahman, K., Heyat, M. B. B., Sumbul, ., Akhtar, F., & Muaad, A. Y. (2022). Role of Inflammation, Oxidative Stress, and Mitochondrial Changes in Premenstrual Psychosomatic Behavioral Symptoms with Anti-Inflammatory, Antioxidant Herbs, and Nutritional Supplements. *Oxidative Medicine and Cellular Longevity*, *2022*, 1-29.

[40] Ferries-Rowe, E., Corey, E., & Archer, J. S. (2020). Primary Dysmenorrhea. *Obstetrics & Gynecology*, *136*(5), 1047-1058.

[41] Iacovides, S., Avidon, I., & Baker, F. C. (2015). What we know about primary q bnm.

[42] Itani, R., Soubra, L., Karout, S., Rahme, D., Karout, L., & Khojah, H. M. J. (2022). Primary Dysmenorrhea: Pathophysiology, Diagnosis, and Treatment Updates. *Korean Journal of Family Medicine*, *43*(2), 101-108.

[43] Martire, F. G., Piccione, E., Exacoustos, C., & Zupi, E. (2023). Endometriosis and Adolescence:The Impact of Dysmenorrhea. *Journal of Clinical Medicine*, *12*(17), 5624.

[44] Stat Pearls (2023). *Dysmenorrhea*. NCBI Bookshelf. NBK560834

[45] Ergonül, P. G., Ergonül, B., & Seçkin, A. K. (2010). Cholesterol content and fatty acid profile of chocolates mostly consumed in Turkey. *CyTA - Journal of Food*, *8*(1), 73–78.

[46] Razola-Díaz, M. C., Aznar-Ramos, M. J., Verardo, V., Melgar-Locatelli, S., Castilla-Ortega, E., & Rodríguez-Pérez, C. (2023). Exploring the Nutritional Composition and Bioactive Compounds in Different Cocoa Powders. *Antioxidants*, *12*(3), 716.

[47] Jalil, A. M. M., & Ismail, A. (2008). Polyphenols in Cocoa and Cocoa Products: Is There a Link between Antioxidant Properties and Health? *Molecules*, *13*(9), 2190–2219.

[48] Melo, T. S., Pires, T. C., Engelmann, J. V. P., Monteiro, A. L. O., Maciel, L. F., & Bispo, E. S. (2020). Evaluation of the content of bioactive compounds in cocoa beans during the fermentation process. *Journal of Food Science and Technology*, *58*(5), 1947–1957.

[49] Quek, R. Y. C., Peh, E. W. Y., & Henry, C. J. (2020). Effects of Cocoa Butter and Cocoa Butter Equivalent in a Chocolate Confectionery on Human Blood Triglycerides, Glucose and Insulin. *Foods*, *9*(4), 455.

[50] Singh, P. K., Khedkar, R. D., & Chandra, S. (2024). Chocolate: An overview of functional potential and recent trends in fortification. *Brazilian Journal of Food Technology*, *27*.

[51] Soares, T. F., & Oliveira, M. B. P. P. (2022). Cocoa By-Products: Characterization of Bioactive Compounds and Beneficial Health Effects. *Molecules*, *27*(5), 1625.

[52] Kababie-Ameo, R., Rabadán-Chávez, G. M., Vázquez-Manjarrez, N., & Gutiérrez-Salmeán, G. (2022). Potential applications of cocoa (Theobroma cacao) on diabetic neuropathy: mini-review. *Frontiers in Bioscience-Landmark*, *27*(2).

[53] Bali, S. (2025). Zesty Pomegranate Bon-Bon Candies: A Tasty Boost For Your Brain And Gut Health.*International Journal of Medical and Pharmaceutical Research*, *6*(6), 461–475.

[54] Maharani, *et al*. Effect of dark chocolate consumption on menstrual pain among late adolescents. 2017. Volume 3, Issue 6. 686–692.

[55] Karakuş Selçuk, Baysal, *et al*. Effect of dark chocolate combined with relaxation therapy on dysmenorrhea pain and anxiety. 2022. Volume 56 102192

[56] Ferina, *et al*. Impact of dark chocolate on dysmenorrhea among adolescents. 2023. Volume 11, Issue 2. 192–195

[57] Fatkurrohman, *et al*. Effectiveness of dark chocolate in reducing dysmenorrhea in school- based adolescents. 2023. Volume 3, Issue 2. 105–111

[58] Okta Viantri, *et al*. Role of dark chocolate in managing dysmenorrhea among adolescents in educational settings. 2025. Volume 17, Issue 1. 48–54

[59] Firnandani S, Achyar K, *et al*. Dark chocolate consumption and its effect on dysmenorrhea: a systematic review. 2025. Volume 13, Issue 1. 47–58

[60] Maharani, *et al*. Effect of dark chocolate consumption on menstrual pain among late adolescents. 2017. Volume 3, Issue 6. 686–692.

[61] Karakuş Selçuk, Baysal, *et al*. Effect of dark chocolate combined with relaxation therapy on dysmenorrhea pain and anxiety. 2022. Volume 56 102192.

[62] Ferina, *et al*. Impact of dark chocolate on dysmenorrhea among adolescents. 2023. Volume 11, Issue 2. 192–195.

[63] Fatkurrohman, *et al*. Effectiveness of dark chocolate in reducing dysmenorrhea in school-based adolescents. 2023. Volume 3, Issue 2. 105–111.

[64] Okta Viantri, *et al*. Role of dark chocolate in managing dysmenorrhea among adolescents in educational settings. 2025. Volume 17, Issue 1. 48–54.

[65] Firnandani S, Achyar K, *et al*. Dark chocolate consumption and its effect on dysmenorrhea: a systematic review. 2025. Volume 13, Issue 1. 47–58.